

# *Allophoma brasiliensis* J.L.V.R. Carvalho, J.D.P. Bezerra & Souza-Motta (Didymellaceae, Pleosporales): first occurrence in an agricultural site from Brazilian Atlantic Forest

Thays G. L. Oliveira<sup>1</sup>, Rejane M. F. Silva<sup>1</sup>, Alexandre R. Machado<sup>1</sup>, Oliane M. C. Magalhães<sup>1</sup>, Antonio F. Costa<sup>2</sup>, Cristina M. Souza-Motta<sup>1</sup>, Gladstone A. Silva<sup>1</sup>

<sup>1</sup> Departamento de Micologia Prof. Chaves Batista, Centro de Biociências, Universidade Federal de Pernambuco, Cidade Universitária, Recife, PE, Brazil

<sup>2</sup> Instituto Agrônomo de Pernambuco, Bongi, Recife, PE, Brazil

Corresponding author: Rejane M. F. Silva (re.biologicas@gmail.com)

**Abstract.** *Allophoma brasiliensis* J.L.V.R. Carvalho, J.D.P. Bezerra & Souza-Motta is reported for the first time as endophyte from leaves of *Capsicum annuum* L. (Solanaceae) isolated in an agricultural site from Brazilian Atlantic Forest. For species determination, morphological characters were analysed along with a multigene analysis performed using the ITS region of the rDNA, the partial large subunit nuclear ribosomal RNA gene (LSU rRNA), the partial second largest subunit of the RNA polymerase II gene subunit (*rpb2*), and the partial  $\beta$ -tubulin gene (*tub2*). A key to *Allophoma* Qian Chen & L. Cai species is provided. This new record contributes to the global fungal diversity, indicating a new geographical distribution, and a new lifestyle.

**Key words.** Bell pepper, endophytic fungi, multigene analysis, *Phoma*-like, taxonomy

**Oliveira TGL, Silva RMF, Machado AR, Magalhães OMC, Costa AF, Souza-Motta CM, Silva GA** (2024) *Allophoma brasiliensis* J.L.V.R. Carvalho, J.D.P. Bezerra & Souza-Motta (Didymellaceae, Pleosporales): first occurrence in an agricultural site from Brazilian Atlantic Forest. Check List 20 (4): 880–888. <https://doi.org/10.15560/20.4.880>

## INTRODUCTION

The genus *Allophoma* Qian Chen & L. Cai was originally introduced by Chen et al. (2015) in a taxonomic review of *Phoma*-like genera within Didymellaceae to accommodate five species originally described as “*Phoma*”: *A. labilis* (Sacc.) Q. Chen & L. Cai (syn. *Phoma labilis*), *A. minor* (Aveskamp, Gruyter & Verkley) Q. Chen & L. Cai (syn. *Phoma minor*), *A. piperis* (Tassi) Q. Chen & L. Cai (syn. *Phoma piperis*), *A. tropica* (R. Schneid. & Boerema) Q. Chen & L. Cai (syn. *Phoma tropica*), *A. zantedeschiae* (Dippen.) Q. Chen & L. Cai (syn. *Phoma zantedeschiae*), and a new species, *A. nicaraguensis* Q. Chen & L. Cai.

*Allophoma* is a monophyletic lineage in Didymellaceae (Chen et al. 2015) and currently comprises 15 species (Hou et al. 2020a; Wijayawardene et al. 2020). *Allophoma tropica* is the type species of the genus. The asexual morphological features of *Allophoma* are represented by the following characters: one conidiomata pycnidial globose to flask-shaped with ostioles; pycnidial wall (2–5-layered) pseudoparenchymatous; conidiogenous cells phialidic, ampulliform to doliiform; and conidia variable in shape and size and mostly guttulate (Chen et al. 2015).

*Allophoma* species have been also isolated from coral (*Acropora formosa* Dana) (Hou et al. 2020b), stems (Hou et al. 2020b), twigs (Chen et al. 2015), and leaves (Marin-Felix et al. 2019) of wide host range as saprobes (de Gruyter et al. 1993; Jayasiri et al. 2019), and/or as causal agents of leaf-spot (de Gruyter et al. 1993; Yuan et al. 2021), dieback, and canker diseases (Babaahmadi et al. 2018). Representatives of this genus have also been found in the air (Chen et al. 2017) and in human-eye lesions (Valenzuela-Lopez et al. 2018).

The Atlantic Forest is one of the richest Neotropical hotspots in terms of diversity, encompassing 17 Brazilian states. However, only 12.4% of forest remnants and natural areas of the original vegetation remain (SOS Mata Atlântica 2017). Degraded areas, including by crop plantations, are remarkable within the heterogeneous ecosystems characteristic of the biome (Barbosa et al. 2022; Wilson et al. 2021), and several fungal species from cultivated areas have been described in the last years (Oliveira et al. 2016; Silva et al. 2019).



Academic editor: Renan Barbosa

Received: 12 December 2023

Accepted: 27 June 2024

Published: 23 July 2024

Copyright © The authors. This is an open-access article distributed under terms of the Creative Commons Attribution License (Attribution 4.0 International – CC BY 4.0)

Analyzing the diversity of endophytic fungi in leaves of *Capsicum annuum* L. (Solanaceae), which is cultivated under the conventional management system, we identified the isolates URM 8563 and URM 8564 as *Phoma*-like based on their morphological characters. After combined multigene sequence analysis of the ITS, LSU rRNA, *tub2*, and *rpb2*, we identify *Allophoma brasiliensis* J.L.V.R. Carvalho, J.D.P. Bezerra & Souza-Motta (Didymellaceae, Pleosporales), and we highlight the occurrence for the first time as endophyte in leaves of *C. annuum* isolated in an agricultural site from Brazilian Atlantic Forest. In addition, we propose a key to identification the species of the genus.

## METHODS


**Collecting and isolation.** Healthy leaf samples of *Capsicum annuum* were collected in 2021 in an agricultural area of conventional cultivation located in the municipality of Chã Grande (08°15'18"S, 035°29'47"W), Pernambuco, Brazil (Figure 1). The agricultural area is inserted within the Brazilian Atlantic Forest biome at an altitude of 488 m. Endophytic fungi were isolated according to the methodology described by Araújo et al. (2002), modified. The leaves were washed in running water. Afterwards, leaf discs were cut with a sterile metal cork punch (6 mm diameter) and disinfested with 70% ethanol for 30 s, in sodium hypochlorite (3% of active chlorine) for 2 min, and finally washed three times in distilled, sterilized water. After isolation and purification, the colonies were kept in water with 10% glycerol for further analysis.

Representative isolates were deposited in the culture collection of the Micoteca URM Profa. Maria Auxiliadora Cavalcanti WCDM 604 at the Universidade Federal de Pernambuco (UFPE), Recife, Brazil.

**Morphological analysis.** Isolates of *Allophoma* were cultured on potato dextrose agar (PDA) and malt extract agar (MEA), and incubated up to 15 days at room temperature in a natural ( $\pm 28^\circ\text{C}$ ) day–night cycle. The isolates sporulated after 15 days, and general culture characteristics were observed (colonies were measured and macroscopic features examined), as well as reproductive structures. Micromorphological characters such as pycnidia (30), conidiogenous cells (three) and conidia (30) were randomly selected and measured. The average of the measurements was used to estimate the size of the structures. The colony colours were described according to Rayner's colour charts (Rayner 1970).

**DNA extraction, PCR amplification, and sequencing.** Fungal biomass was obtained from colonies grown on PDA plates at room temperature for up to 15 days. Genomic DNA extraction was performed, with the material previously ground, as described by Oliveira et al. (2016). The primers ITS1/ITS4 (White et al. 1990), LR0R/LR5 (Vilgalys and Hester 1990; Vilgalys and Sun 1994), fRPB2-5F2/fRPB2-7cR (Liu et al. 1999; Sung et al. 2007), and Bt2a/Bt2b (Glass and Donaldson 1995) were used to amplify the ITS region of the rRNA (ITS), the partial large subunit nuclear ribosomal RNA gene (LSU rRNA), the partial second largest subunit of the RNA polymerase II gene (*rpb2*) subunit, and the partial  $\beta$ -tubulin gene (*tub2*), respectively. The PCR conditions for ITS and LSU rRNA were an initial denaturation at  $95^\circ\text{C}$  for 3 min, followed by 39 cycles of

**Figure 1.** Map of the occurrence of *Allophoma brasiliensis* J.L.V.R. Carvalho, J.D.P. Bezerra & Souza-Motta for the first time as an endophyte in *Capsicum annuum* leaves on an agricultural site in the Brazilian Atlantic Forest.



denaturation at 94 °C for 45 s, annealing at 58 °C for 1 min, and extension at 72 °C for 1 min, with a final extension step at 72 °C for 7 min. The PCR conditions for *rpb2* was an initial denaturation at 95 °C for 5 min, followed by 40 cycles of denaturation at 95 °C for 1 min, annealing at 57 °C for 2 min and extension at 72 °C for 90 s, with a final extension step at 72 °C for 10 min. The PCR conditions for *tub2* was an initial denaturation at 94 °C for 5 min, followed by 35 cycles of denaturation at 94 °C for 45 s, annealing at 55 °C for 45 s and extension at 72 °C for 1 min, with a final extension step at 72 °C for 7 min. The amplification reactions were carried out in a final volume of 25 µl. All products obtained from the amplification were purified using the enzyme Exonuclease I (Thermo Fisher Scientific) and Shrimp Alkaline Phosphatase (Cellco) and sequenced on the Multiuser Platform for Sequencing and Gene Expression of the UFPE, Centro de Biociências.

**Sequence alignment and phylogenetic analysis.** For the final molecular characterization of the species of *Allophoma*, four different markers were used: ITS, LSU rRNA, *rpb2*, and *tub2*. The dataset was constructed with the sequences of our isolates together with sequences of type material and representative isolates, totaling 28 sequences of the 15 accepted species of *Allophoma*. *Stagonosporopsis loticola* (Died.) Aveskamp, Gruyter & Verkley (CBS 562.81) was used as the outgroup. The GenBank accession numbers of all sequences used in this study and the newly obtained sequences are shown in Appendix Table A1. Sequences were aligned using the online tool MAFFT v. 7 (Katoh and Standley 2013) and manually edited with the MEGA v. 7 software (Kumar et al. 2016).

The concatenated four-locus datasets were analyzed by maximum likelihood (ML) and bayesian inference (BI) using the RAxML-HPC BlackBox v. 8.2.12 (Stamatakis 2014) and MrBayes on XSEDE v. 3.2.7a (Ronquist et al. 2012), respectively, at the online server of the Cipres Science gateway portal (Miller et al. 2010). ML analyzes were performed with 1000 bootstrap replications using default parameters and the GTR + G + I model test. For BI analyses, the best nucleotide substitution model was estimated by MrModelTest v. 2.3 (Nylander 2004) for each gene region separately (ITS = HKY + I, LSU = GTR + I, *rpb2* = SYM + G, and *tub2* = GTR + G). BI analyzes were performed with two Markov Chain Monte Carlo (MCMC),  $1 \times 10^7$  generations with a burn value of 25%, and trees were sampled every 1000 generations. In the nodes, BI posterior probability (BPP) and ML bootstrap (ML–BS) values equal to or greater than 0.95 and 70 %, respectively, are presented. The ultimate phylogenetic tree was visualized in FigTree v. 1.4.3 (<http://tree.bio.ed.ac.uk/software/figtree/>). The concatenated alignment was deposited in the TreeBASE (study ID 29859).

## RESULTS

***Allophoma brasiliensis*** J.L.V.R. Carvalho, J.D.P. Bezerra & Souza-Motta, 2022;  
Brazilian Journal of Microbiology, emend.  
Figure 2A–J

**Materials examined.** BRAZIL – PERNAMBUCO • Chã Grande, Rancho Bela Vista; 08°15'18"S, 035°29'47"W; 488 m alt; 29.I.2021; T.G.L. Oliveira leg.; as endophytic fungi in leaves of *Capsicum annuum*; URM 8563 • ibid.; URM 8564.

**Distribution.** The new record was found as endophyte in leaves of *C. annuum* in an agricultural area inserted in Brazilian Atlantic Forest.

**Identification** (colony characters, 28 °C, 15 days). PDA: colony regular and white margins, greenish olivaceous, black pycnidia forming a concentric ring, aerial mycelium sparse and whitish; opaque to greenish olivaceous reverse, white near margins. MEA: colony opaque to greenish olivaceous, covered by whitish flaky aerial mycelium, irregular margin, light brown to white near the colony margin; opaque to greenish olivaceous reverse.

**Colony diameters**, 15 days, in cm. PDA 28 °C 5.8; MEA 28 °C 5.8.

**Micromorphology** (28 °C, 15 days). PDA: *conidiomata* pycnidial, produced on the agar surface or semi-immersed, mostly isolated, some grouped, globose or subglobose to flask-shaped, pale brown, some dark brown, thin-walled, glabrous, ostiolate, (60–)108–192(–216) × (60–)120–180(–192) µm. *Ostiole* single, slightly papillate. *Conidiogenous cells* phialidic, hyaline, globose to ampulliform, smooth, slightly papillate, 3.5–5(–6.3) × 2.2–3.7(–5.2) µm. *Conidia* oblong with both ends rounded, smooth- and thin-walled, hyaline, aseptate, without guttules, 3.1–4.7(–6.6) × 1.3–2.2 µm. *Conidial matrix* cream. *Chlamydospores* multicellular, mostly intercalary in chain, smooth, dark brown.

**Comments.** According to phylogenetic analyses, strains URM 8563 and URM 8564 grouped into the same highly supported clade (BPP = 0.95 and ML–BS = 100) together with *A. brasiliensis* ex-type strain (URM 8453) and *A. brasiliensis* URM 8454, isolated from bat flies (*Trichobius* sp. Gervais) in Catimbau National Park, Brazil (Figure 3). Some morphological differences between the endophytes (URM 8563 and URM 8564), isolated from leaves of *C. annuum*, and *A. brasiliensis* ex-type strain (URM 8453) were observed. We




**Figure 2.** *Allophoma brasiliensis* URM 8563. **A.** Detail of colony on PDA after 15 days. **B.** Detail of colony on MEA after 15 days. **C, D.** Conidiomata pycnidial. **E–G.** Conidiogenous cells and conidia. **H, I.** Conidia. **J.** Chlamydospores. Scale bars: C, D = 100  $\mu$ m; E–J = 10  $\mu$ m.

have observed in our specimens pycnidia larger, and conidiogenous cells smaller than ex-type strain (URM 8453). Chlamydospores and conidia without guttules were present in our strains and were not observed in ex-type strain (URM 8453). Furthermore, *A. brasiliensis* (URM 8563 and URM 8564) is reported for the first time as an endophytic fungus.

#### Key to the *Allophoma* species

- |     |                                                                                                      |                         |
|-----|------------------------------------------------------------------------------------------------------|-------------------------|
| 1a  | Conidiomata pycnidial (sub-)globose to ellipsoidal or flask-shaped                                   | 2                       |
| 1b  | Conidiomata pycnidial ovoid                                                                          | 3                       |
| 2a  | Conidiomata pycnidial up to 300 $\mu$ m long                                                         | 4                       |
| 2b  | Conidiomata pycnidial 301–635 $\mu$ m long                                                           | 5                       |
| 3a  | Conidiomata pycnidial 120–210 $\times$ 90–140 $\mu$ m                                                | <i>A. cylindrispora</i> |
| 3b  | Conidiomata pycnidial 70–90 $\times$ 68–85 $\mu$ m                                                   | <i>A. siamensis</i>     |
| 4a  | Ostiole 1 single, slightly papillate, papillate, or non-papillate                                    | 6                       |
| 4b  | Ostioles 1–3 slightly papillate, papillate or sometimes non-papillate                                | 7                       |
| 5a  | Ostioles 1–3 slightly papillate, papillate, or non-papillate                                         | 8                       |
| 5b  | Ostioles 1–5 slightly papillate, papillate, or non-papillate                                         | 9                       |
| 6a  | Conidia oblong to ellipsoidal, hyaline, aseptate, with polar guttules                                | 10                      |
| 6b  | Conidia oval, oblong or ellipsoidal, hyaline, aseptate, eguttulate                                   | 11                      |
| 7a  | Conidia oblong to cylindrical, hyaline, aseptate, with minutes guttules                              | <i>A. thunbergiae</i>   |
| 7b  | Conidia oblong to ellipsoidal, hyaline to pale brown, 0- or 1-septate, majority 1-septate, guttulate |                         |
|     |                                                                                                      | <i>A. hayatii</i>       |
| 8a  | Conidia oblong, hyaline, aseptate, with 1 or 2 medium-sized polar guttules                           | <i>A. anatae</i>        |
| 8b  | Conidia oblong, hyaline, aseptate, with 2 large polar guttules                                       | <i>A. alba</i>          |
| 8c  | Conidia oblong to cylindrical, hyaline, aseptate, with 2 distinct pale-green polar guttules          |                         |
|     |                                                                                                      | <i>A. oligotrophica</i> |
| 9a  | Conidia ellipsoidal to ovoid or slightly allantoid, hyaline, aseptate, (0–)1–3(–4) minute guttules   | <i>A. minor</i>         |
| 9b  | Conidia ellipsoidal to ovoid or oblong, hyaline, aseptate, 1 or 2 guttules                           | 12                      |
| 10a | Conidial matrix whitish to pale luteous or ochraceous                                                | 13                      |
| 10b | Conidial matrix not recorded                                                                         | <i>A. piperis</i>       |
| 11a | Conidial matrix greyish and presence of chlamydospores in the mycelium                               | <i>A. zantedeschiae</i> |
| 11b | Conidial matrix cream and presence of chlamydospores in the mycelium                                 | <i>A. brasiliensis</i>  |
| 12a | Conidial matrix white-yellowish                                                                      | <i>A. tropica</i>       |

- 12b Conidial matrix cream ..... *A. pterospermicola*  
 13a Negative NaOH spot-test on oatmeal agar ..... *A. nicaraguensis*  
 13b Positive NaOH spot-test on oatmeal agar ..... *A. labilis*


## DISCUSSION

Studying the diversity of endophytic fungi present in bell pepper leaves, we isolated the endophytes URM 8563 and URM 8564, whose morphological characteristics belonged to the *Phoma*-like group. Based on combined multigene sequence analysis of the ITS, LSU rRNA, *rpb2*, and *tub2*, our phylogenetic analyses indicated that our specimens belong to *Allophoma brasiliensis*, which is reported here for the first time as endophyte.

Other species of the genus *Allophoma* have been reported as endophytes elsewhere in the world, including in the stems and wilted leaves of *Ageratina adenophora* (Spreng.) R. M. King & H. Rob. in Yunnan Province, southwestern China (Fang et al. 2019) and in leaves of *Rhizophora apiculata* Blume in Phetchaburi Province, Thailand (Doilom et al. 2017). Members of *Allophoma* are characterized by aseptate conidia that vary both in shape and size, are hyaline and mostly guttulate, and have globose and ostiolate pycnidia (Chen et al. 2015).

The endophytic isolates obtained in this study (URM 8563 and URM 8564) are similar to the description of *Allophoma brasiliensis* culture ex-type URM 8453 proposed by Carvalho et al. (2022), which was isolated from bat flies (*Trichobius* sp.) from Catimbau National Park. This park is within the Caatinga domain and encompasses parts of the municipalities of Buíque, Tupanatinga, and Ibimirim in the state of Pernambuco. Carvalho et al. (2022) used macromorphological characters from colonies in OA, PDA, and MEA cultivated at 27 °C for 7 days in the dark and micromorphological characters such as conidiomata pycnidial, conidiomatal wall, conidiogenous cells, and conidia for identification and confirmation of the species. Multigene analysis (ITS, LSU rRNA, *rpb2*, and *tub2*) demonstrated high identity and phylogenetic relationship with our isolates (Figure 3).

**Figure 3.** Phylogenetic tree generated from Bayesian inference analysis based on combined *tub2*, *rpb2*, ITS, and LSU rRNA sequences. Sequences from type species and the reference isolates are indicated in bold. The sequence obtained in this study is annotated in red. Bayesian posterior probabilities (above 0.95) and maximum likelihood bootstrap (above 70%) values are shown near nodes. The tree is rooted with *Stagonosporopsis loticola* (CBS 562.81).



The differences in the lifestyle and environment of the isolation for the strains of this fungi possibly are responsible for some variability in morphological characters found. Versatility in the morphological and ecophysiological characteristics of fungi may occur as a form of adaptation to nutrient acquisition, stress tolerance, and interactions with other organisms, in addition to mediating the biogeography of these organisms (Bahram and Netherway 2022). We observed chlamydospores in our specimens, although this was not reported by Carvalho et al. (2022). These authors also have found guttulated conidia, but these structures were not observed by us. The pycnidia found in our isolates were larger than those observed by Carvalho et al. (2022), while the conidiogenous cells from URM 8563 and URM 8564 are smaller than those in the species' original description.

*Allophoma brasiliensis* (URM 8563 and URM 8564) was found in an agricultural area in the municipality of Chã Grande as endophyte from healthy leaves of Bell Pepper, *Capsicum annuum*, and *A. brasiliensis* species is reported here for the first time in an agricultural site in Brazilian Atlantic Forest. Our study is an important addition to knowledge of the global fungal diversity, as it adds a new geographical occurrence, expands the known geographical distribution, includes a different, previously unreported biome, and reports a new ecological lifestyle for *A. brasiliensis*.

## ACKNOWLEDGEMENTS

We are grateful to the Hortalíça Center J.C. Alves e CIA Ltda for allowing access to cultivation areas to carry out the collections. This work was supported through a research grant from the Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco (IBPG-0823-2.03/20). This research received financial support from the Universidade Federal de Pernambuco (Edital PROPG no. 06/2022). Cristina M. de Souza-Motta and Gladstone A. Silva thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the research grants (processes 311187/2022-6 and 312606/2022-2).

## ADDITIONAL INFORMATION

### Conflict of interest

The authors declare that no competing interests exist.

### Ethical statement

No ethical statement is reported.

### Funding

This study was financially supported by Universidade Federal de Pernambuco (Edital PROPG no. 06/2022), and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the research grants (processes 311187/2022-6 and 312606/2022-2).

### Author contributions

Conceptualization: TGLO, RMFS, ARM, OMCM, CMSM, AFC, GAS. Data curation: TGLO, RMFS, ARM, OMCM, CMSM, AFC, GAS. Methodology: TGLO, RMFS, GAS. Formal analysis: TGLO, RMFS, ARM, OMCM, AFC, GAS. Visualization: TGLO, RMFS, ARM, OMCM, AFC, GAS. Funding acquisition: GAS. Resources: GAS. Supervision: GAS, ARM, OMCM, TGLO. Project administration: GAS, TGLO. Writing – original draft: TGLO, GAS. Writing – review and editing: TGLO, GAS.

### Author ORCID iDs

Thays G. L. Oliveira  <https://orcid.org/0000-0001-5415-3916>

Rejane M. F. Silva  <https://orcid.org/0000-0002-5024-0286>

Alexandre R. Machado  <https://orcid.org/0000-0001-7440-3097>

Oliane M. C. Magalhães  <https://orcid.org/0000-0001-7275-5304>

Antonio F. Costa  <https://orcid.org/0000-0001-9866-3504>

Cristina M. Souza-Motta  <https://orcid.org/0000-0002-0964-8271>

Gladstone A. Silva  <https://orcid.org/0000-0002-5652-7800>

### Data availability

All data that support the findings of this study are available in the main text and the Appendix.

## REFERENCES

- Araújo WL, Lima AOS, Azevedo JL, Marcon J, Sobral JK, Lacava PT (2002) Manual: isolamento de microrganismos endofíticos. Centro Acadêmico Luiz de Queiroz, Piracicaba, Brasil, 86 pp.
- Babaahmadi G, Mehrabi-Koushki M, Hayati J (2018) *Allophoma hayatii* sp. nov., an undescribed pathogenic fungus causing dieback of *Lantana camara* in Iran. Mycological Progress 17: 365–379. <https://doi.org/10.1007/s11557-017-1360-7>

- Bahram M, Netherway T** (2022) Fungi as mediators linking organisms and ecosystems. *FEMS Microbiology Reviews* 46 (2): fuab058. <https://doi.org/10.1093/femsre/fuab058>
- Barbosa RN, Santos JEF, Bezerra JDP, Istel Ł, Houbraken J, Oliveira NT, Souza-Motta CM** (2022) Brazilian Atlantic Forest and Pampa biomes in the spotlight: an overview of *Aspergillus*, *Penicillium*, and *Talaromyces* (Eurotiales) species and the description of *Penicillium nordestinense* sp. nov. *Acta Botanica Brasilica* 36: e2021abb0390. <https://doi.org/10.1590/0102-33062021abb0390>
- Carvalho JLVR, Lima JMS, Barbier E, Bernard E, Bezerra JDP, Souza-Motta CM** (2022) Ticket to ride: fungi from bat ectoparasites in a tropical cave and the description of two new species. *Brazilian Journal of Microbiology* 53 (4): 2077–2091. <https://doi.org/10.1007/s42770-022-00841-y>
- Chen Q, Hou LW, Duan WJ, Crous PW, Cai L** (2017) *Didymellaceae* revisited. *Studies in Mycology* 87 (1): 105–159. <https://doi.org/10.1016/j.simyco.2017.06.002>
- Chen Q, Jiang JR, Zhang GZ, Cai L, Crous PW** (2015) Resolving the *Phoma* enigma. *Studies in Mycology* 82: 137–217. <https://doi.org/10.1016/j.simyco.2015.10.003>
- De Gruyter J, Noordeloos ME, Boerema GH** (1993) Contributions towards a monograph of *Phoma* (Coelomycetes)—I. 2. Section *Phoma*: additional taxa with very small conidia and taxa with conidia up to 7 µm long. *Persoonia—Molecular Phylogeny and Evolution of Fungi* 15 (3): 369–400.
- Doilom M, Manawasinghe IS, Jeewon R, Jayawardena RS, Tibpromma S, Hongsanan S, Meepol W, Lumyong S, Jones EBG, Hyde KD** (2017) Can ITS sequence data identify fungal endophytes from cultures? A case study from *Rhizophora apiculata*. *Mycosphere* 8 (10): 1869–1892. <https://doi.org/10.5943/mycosphere/8/10/11>
- Fang K, Miao Y-F, Chen L, Zhou J, Yang ZP, Dong XF, Zhang HB** (2019) Tissue-specific and geographical variation in endophytic fungi of *Ageratina adenophora* and fungal associations with the environment. *Frontiers in Microbiology* 10: 2919. <https://doi.org/10.3389/fmicb.2019.02919>
- Glass NL, Donaldson GC** (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Applied and Environmental Microbiology* 61 (4): 1323–1330. <https://doi.org/10.1128/aem.61.4.1323-1330.1995>
- Hou L, Hernández-Restrepo M, Groenewald JZ, Cai L, Crous PW** (2020a) Citizen science project reveals high diversity in *Didymellaceae* (Pleosporales, Dothideomycetes). *Mycosphere* 65: 49–99. <https://doi.org/10.3897/mycokeys.65.47704>
- Hou LW, Groenewald JZ, Pfenning LH, Yarden O, Crous PW, Cai L** (2020b) The *Phoma*-like dilemma. *Studies in Mycology* 96: 309–396. <https://doi.org/10.1016/j.simyco.2020.05.001>
- Jayasiri SC, Hyde KD, Jones EBG, McKenzie EHC, Jeewon R, Phillips AJL, Bhat DJ, Wanasinghe DN, Liu JK, Lu YZ, Kang JC, Xu J, Karunarathna SC** (2019) Diversity, morphology and molecular phylogeny of Dothideomycetes on decaying wild seed pods and fruits. *Mycosphere* 10 (1): 1–186. <https://doi.org/10.5943/mycosphere/10/1/1>
- Katoh K, Standley DM** (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution* 30: 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kumar S, Stecher G, Tamura K** (2016) MEGA7: Molecular Evolutionary Genetics Analysis version 7.0 for Bigger Datasets. *Molecular Biology and Evolution* 33 (7): 1870–1874. <https://doi.org/10.1093/molbev/msw054>
- Liu YJ, Whelen S, Hall BD** (1999) Phylogenetic relationships among ascomycetes: evidence from an RNA polymerase II subunit. *Molecular Biology and Evolution* 16 (12): 1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>
- Marin-Felix Y, Hernández-Restrepo M, Iturrieta-González I, García D, Gené J, Groenewald JZ, Cai L, Chen Q, Quaedvlieg W, Schumacher RK, Taylor PWJ, Ambers C, Bonthond G, Edwards J, Krueger-Hadfield SA, Luangsa-Ard JJ, Morton L, Moslemi A, Sandoval-Denis M, Tan YP, Thangavel R, Vaghefi N, Cheewangkoon R, Crous PW** (2019) Genera of phytopathogenic fungi: GOPHY 3. *Studies in Mycology* 94: 1–124. <https://doi.org/10.1016/j.simyco.2019.05.001>
- Miller MA, Pfeiffer W, Schwartz T** (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. *Gateway Computing Environments Workshop (GCE)* 1–8. <https://doi.org/10.1109/gce.2010.5676129>
- Nylander JAA** (2004) MrModeltest Version 2. Program Distributed by the Author. Evolutionary Biology Centre, Uppsala University.
- Oliveira RJV, Bezerra JL, Lima TEF, Silva GA, Cavalcanti MAQ** (2016) *Phaeosphaeria nodulispora*, a new endophytic coelomycete isolated from tropical palm (*Cocos nucifera*) in Brazil. *Nova Hedwigia* 103 (1): 185–192. [https://doi.org/10.1127/nova\\_hedwigia/2016/0343](https://doi.org/10.1127/nova_hedwigia/2016/0343)
- Rayner RW** (1970) A mycological colour chart. CMI and British Mycological Society, Kew, UK.
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP** (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61 (3): 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Silva RM, Oliveira RJ, Bezerra JD, Bezerra JL, Souza-Motta CM, Silva GA** (2019) *Bifusisporrella sorghi* gen. et sp. nov. (Magnaporthaceae) to accommodate an endophytic fungus from Brazil. *Mycological Progress* 18: 847–854. <https://doi.org/10.1007/s11557-019-01494-2>
- SOS Mata Atlântica** (2017) Relatório Anual de atividades 2017. [https://www.sosma.org.br/wp-content/uploads/2018/10/AF\\_RA\\_SOSMA\\_2017\\_web.pdf](https://www.sosma.org.br/wp-content/uploads/2018/10/AF_RA_SOSMA_2017_web.pdf). Accessed on: 2023-06-21.
- Stamatakis A** (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30 (9): 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>

- Sung GH, Sung JM, Hywel-Jones NL, Spatafora JW** (2007) A multi-gene phylogeny of Clavicipitaceae (Ascomycota, Fungi): identification of localized incongruence using a combinational bootstrap approach. *Molecular Phylogenetics and Evolution* 44 (3): 1204–1223. <https://doi.org/10.1016/j.ympev.2007.03.011>
- Valenzuela-Lopez N, Cano-Lira JF, Guarro J, Sutton DA, Wiederhold N, Crous PW, Stchigel AM** (2018) Coelomycetous Dothideomycetes with emphasis on the families Cucurbitariaceae and Didymellaceae. *Studies in Mycology* 90: 1–69. <https://doi.org/10.1016/j.simyco.2017.11.003>
- Vilgalys R, Hester M** (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172 (8): 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- Vilgalys R, Sun BL** (1994) Ancient and recent patterns of geographic speciation in the oyster mushroom *Pleurotus* revealed by phylogenetic analysis of ribosomal DNA sequences. *Proceedings of the National Academy of Sciences of the United States of America* 91 (10): 4599–4603. <https://doi.org/10.1073/pnas.91.10.4599>
- White TJ, Bruns T, Lee S, Taylor J** (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds.) *PCR protocols: a guide to methods and applications*. Academic Press, New York, USA, 315–322.
- Wijayawardene NN, Hyde KD, Al-Ani LKT, Tedersoo L, Haelewaters D, Rajeshkumar KC, Zhao RL, Aptroot A, Leontyev DV, Saxena RK, et al.** (2020) Outline of Fungi and fungus-like taxa. *Mycosphere* 11 (1): 1060–1456. <https://doi.org/10.5943/mycosphere/11/1/8>
- Wilson OJ, Mayle FE, Walters RJ, Lingner DV, Vibrans AC** (2021) Floristic change in Brazil's Southern Atlantic Forest biodiversity hotspot: from the last glacial maximum to the late 21st century. *Quaternary Science Reviews* 264: 107005. <https://doi.org/10.1016/j.quascirev.2021.107005>
- Yuan J, Zeng X-Y, Geng K, Wijayawardene NN, Bhat JD, Wu SP, Wang Y, Yang ZF** (2021) *Allophoma* species (Pleosporales: Didymellaceae) associated with *Thunbergia grandiflora* in Guangxi Province, China. *Biodiversity Data Journal* 9: e63643. <https://doi.org/10.3897/bdj.9.e63643>

APPENDIX

**Table A1.** Isolates used in this study, including the different markers with respective NCBI GenBank accessions numbers, and the culture code. GenBank accessions numbers in bold were generated in this work.

| Species                          | Culture code*                       | GenBank accession numbers |                 |                 |                 | References              |
|----------------------------------|-------------------------------------|---------------------------|-----------------|-----------------|-----------------|-------------------------|
|                                  |                                     | ITS                       | tub2            | rpb2            | LSU             |                         |
| <i>Allophoma alba</i>            | CBS 120422 <sup>†</sup>             | MN973469                  | MT005568        | MT018044        | MN943671        | Hou et al. 2020b        |
| <i>Allophoma anataiae</i>        | CBS 124673 <sup>†</sup>             | MN973472                  | MT005571        | MT018048        | MN943674        | Hou et al. 2020b        |
| <i>Allophoma brasiliensis</i>    | URM 8453 <sup>†</sup>               | OM692214                  | ON715830        | ON715834        | ON678611        | Carvalho et al. 2022    |
|                                  | URM 8454                            | OM692215                  | ON715831        | ON715835        | ON678612        | Carvalho et al. 2022    |
|                                  | <b>URM 8563; PC74</b>               | <b>OP304832</b>           | <b>OP311925</b> | <b>OP311927</b> | <b>OP304851</b> | <b>This study</b>       |
|                                  | <b>URM 8564; PC75</b>               | <b>OP304833</b>           | <b>OP311926</b> | <b>OP311928</b> | <b>OP304852</b> | <b>This study</b>       |
| <i>Allophoma cylindrispora</i>   | CBS 142453 <sup>†</sup>             | LT592920                  | LT592989        | LT593058        | LN907376        | Hou et al. 2020b        |
| <i>Allophoma hayatii</i>         | CBS 142859 <sup>†</sup>             | KY684812                  | KY684816        | MF095108        | KY684814        | Hou et al. 2020b        |
|                                  | CBS 142860                          | KY684813                  | KY684817        | MF095109        | KY684815        | Hou et al. 2020b        |
| <i>Allophoma labilis</i>         | CBS 124.93; PD 87/269               | GU237765                  | GU237619        | KT389552        | GU238091        | Hou et al. 2020b        |
|                                  | CBS 479.93; PD 70/93 <sup>R</sup>   | GU237868                  | GU237620        | MN983277        | GU238092        | Hou et al. 2020b        |
| <i>Allophoma minor</i>           | CBS 325.82 <sup>†</sup>             | GU237831                  | GU237632        | KT389553        | GU238107        | Hou et al. 2020b        |
|                                  | CBS 315.83 <sup>R</sup>             | GU237826                  | GU237631        | MN983279        | GU238106        | Hou et al. 2020b        |
| <i>Allophoma nicaraguensis</i>   | CBS 506.91; IMI 215229 <sup>†</sup> | GU237876                  | GU237596        | KT389551        | GU238058        | Hou et al. 2020b        |
|                                  | CBS 497.91; PD 79/209               | GU237870                  | GU237597        | —               | GU238059        | Hou et al. 2020b        |
| <i>Allophoma oligotrophica</i>   | CGMCC 3.18114; LC 6245 <sup>†</sup> | KY742040                  | KY742282        | KY742128        | KY742194        | Hou et al. 2020b        |
|                                  | CGMCC 3.18115; LC 6246              | KY742041                  | KY742283        | KY742129        | KY742195        | Hou et al. 2020b        |
| <i>Allophoma piperis</i>         | CBS 268.93; PD 88/720 <sup>†</sup>  | GU237816                  | GU237644        | KT389554        | GU238129        | Hou et al. 2020b        |
|                                  | PD 90/2011                          | GU237921                  | GU237645        | MT018045        | GU238130        | Hou et al. 2020b        |
| <i>Allophoma pterospermicola</i> | CGMCC 3.19245 <sup>†</sup>          | MK088573                  | MK088594        | MK088587        | MK088580        | Marin-Feliz et al. 2019 |
|                                  | LC12181                             | MK088569                  | MK088590        | MK088583        | MK088576        | Marin-Feliz et al. 2019 |
| <i>Allophoma siamensis</i>       | MFLUCC 17-2422 <sup>†</sup>         | MK347742                  | MK412867        | MK434912        | MK347959        | Hou et al. 2020b        |
| <i>Allophoma thunbergiae</i>     | GUCC 2070.7 <sup>†</sup>            | MW036298                  | MW116823        | MW116819        | MW040201        | Yuan et al. 2021        |
| <i>Allophoma tropica</i>         | CBS 436.75; DSM 63365 <sup>†</sup>  | GU237864                  | GU237663        | KT389556        | GU238149        | Hou et al. 2020b        |
|                                  | CBS 537.66; PD 65/27                | MN973468                  | MT005567        | MT018043        | MN943670        | Hou et al. 2020b        |
| <i>Allophoma zantedeschiae</i>   | CBS 131.93; PD 69/140               | FJ427084                  | FJ427188        | KT389557        | GU238159        | Hou et al. 2020b        |
|                                  | CBS 229.32                          | KT389473                  | KT389767        | KT389558        | KT389690        | Hou et al. 2020b        |
| <i>Stagonosporopsis loticola</i> | CBS 562.81; PDDCC 6884 <sup>†</sup> | GU237890                  | GU237697        | KT389684        | GU238192        | Hou et al. 2020b        |

<sup>R</sup>Reference isolates.

<sup>†</sup>Ex-type strain.

\*CBS: Westerdijk Fungal Biodiversity Institute (formerly CBS KNAW), Utrecht, The Netherlands; CGMCC: China General Microbiological Culture Collection, Beijing, China; CPC: Culture collection of Pedro Crous, housed at CBS; DSM: Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany; GUCC: Culture Collection of the Department of Plant Pathology, Agriculture College, Guizhou University; IMI: International Mycological Institute, CABI-Bioscience, Egham, BAKEHAM Lane, UK; LC: LeiCai, corresponding author’s personal collection deposited in laboratory, housed at CAS, China; MFLUCC: Mae Fah Luang University Culture Collection, Chiang Rai, Thailand; PD: Plant Protection Service, Wageningen, the Netherlands; PDDCC: Plant Diseases Division Culture Collection, Auckland, New Zealand; URM: Culture collection Prof. Maria Auxiliadora Cavalcanti, Recife, Brazil.